

Letters

Thyroid Nodule Characterization Using Combined Fine-Needle Aspiration and ^{99m}Tc -Sestamibi Scintigraphy Strategy

In the February 2016 issue of the *AJR*, Yerubandi et al. [1] wrote an interesting article on the frequency of thyroid nodules observed incidentally on non-FDG PET nuclear medicine imaging modalities, including ^{99m}Tc -sestamibi radiotracer, the malignancy rate of such nodules, and predictors of malignancy.

To further expand on this point, it has been shown that fine-needle aspiration (FNA) combined with ^{99m}Tc -sestamibi scintigraphy for evaluating cold thyroid nodules larger than 1 cm in diameter is potentially cost-effective in the management of solitary or dominant thyroid nodules [2]. Sestamibi-based strategies have a lower cost per patient and lower cost per cancer diagnosed and are associated with a low radiation burden [2].

We have observed several cases of discrepancies between FNA cytology and ^{99m}Tc -sestamibi scintigraphy results in our practice, which has evaluated more than 1000 cases to date [3, 4] (Fig. 1). Such cases illustrate the potential complexity of diagnosing thyroid

nodules, and strategies combining FNA with ^{99m}Tc -sestamibi scintigraphy may also end in diagnostic dilemmas.

The discrepancies and dilemmas described here illustrate the need for a pragmatic approach to training medical practitioners so that they may adapt their treatment strategies for patients with cold thyroid nodules.

Iraj Nabipour

Bushehr Medical University Hospital,
Bushehr University of Medical Sciences,
Bushehr, Iran

Mohammadreza Kalantarhormozi
Persian Gulf Tropical Medicine
Research Center,
Bushehr University of Medical Sciences,
Bushehr, Iran

Majid Assadi

Persian Gulf Nuclear Medicine
Research Center,
Bushehr University of Medical Sciences,
Bushehr Medical University Hospital,
Bushehr, Iran

DOI:10.2214/AJR.16.16163

WEB—This is a web exclusive article.

References

1. Yerubandi V, Chin BB, Sosa JA, Hoang JK. Incidental thyroid nodules at non-FDG PET nuclear medicine imaging: evaluation of prevalence and malignancy rate. *AJR* 2016; 206:420–425
2. Wale A, Miles KA, Young B, et al. Combined ^{99m}Tc -methoxyisobutylisonitrile scintigraphy and fine-needle aspiration cytology offers an accurate and potentially cost-effective investigative strategy for the assessment of solitary or dominant thyroid nodules. *Eur J Nucl Med Mol Imaging* 2014; 41:105–115
3. Riazi A, Kalantarhormozi MR, Nabipour I, et al. Technetium-99m methoxyisobutylisonitrile scintigraphy in the assessment of cold thyroid nodules: is it time to change the approach to the management of cold thyroid nodules? *Nucl Med Commun* 2014; 35:51–57
4. Riazi A, Kalantarhormozi M, Nabipour I, Ostovar A, Javadi H, Assadi M. Comments on Wale et al.: Combined ^{99m}Tc -methoxyisobutylisonitrile scintigraphy and fine-needle aspiration cytology offers an accurate and potentially cost-effective investigative strategy for the assessment of solitary or dominant thyroid nodules. (letter) *Eur J Nucl Med Mol Imaging* 2014; 41:575–576




Fig. 1—39-year-old woman with cold thyroid nodule. Scintigraphy with ^{99m}Tc -pertechnetate (left) shows nodule on left lobe of thyroid gland. Nodule shows significant radiotracer retention with poor washout (arrow), which is also seen on sestamibi scan (right). Fine-needle aspiration cytology findings showed nodular goiter, but histopathologic examination showed medullary thyroid carcinoma 4 mm in diameter.